

On geodesic mappings of manifolds with affine connection

Josef Mikeš and Irena Hinterleitner *

Abstract

In this paper we prove that all manifolds with affine connection are globally projectively equivalent to some space with equiaffine connection (equiaffine manifold). These manifolds are characterised by a symmetric Ricci tensor.

1 Introduction

Diffeomorphisms between two spaces A_n and $\bar{A}_n$ with affine connection are called *geodesic*, if any geodesic of A_n is mapped to a geodesic of $\bar{A}_n$, see for example [1] – [15], etc.

Affine (or *trivial geodesic*) *mappings* are mappings which preserve canonical parameters of geodesics. Many papers are devoted to the metrizability or projective metrizability of spaces with affine connection. Under metrizability of a space A_n we understand the existence of a metric g , which generates the affine connection ∇ , such that ∇ is the Levi-Civita connection of g (for which $\nabla g = 0$), see [1, 2, 3, 10]. On the other hand, the problem of metrizability, respectively projective metrizability, is equivalent to that of affine, respectively geodesic, mappings of a space A_n with affine connection onto (pseudo-) Riemannian spaces.

Equiaffine spaces, characterised by the symmetry of the Ricci tensor, are spaces in which the volume of an n dimensional parallelepiped is invariant under parallel transport, play an important role in the theory of geodesic mappings. J. Mikeš and V.E. Berezovski [7, 10] found fundamental equations of geodesic mappings of equiaffine spaces onto (pseudo-) Riemannian spaces in the form of systems of linear partial differential equations of Cauchy type in terms of covariant derivatives. These results were used for further studies by M.G. Eastwood and V. Matveev [1].

*Supported by the Council of the Czech Government MSM 6198959214.

In the paper [9] by I. Hinterleitner, J. Mikeš and V.A. Kiosak it was proved that any space with affine connection is locally projective equiaffine. We show that these properties hold globally, i.e. any arbitrary manifold with affine connection globally admits geodesic mappings onto some equiaffine manifold.

For this reason the solution of the problem of the projective metrizability of a space A_n (or equivalently of geodesic mappings of A_n onto (pseudo-) Riemannian manifolds $\bar{V}_n$) can be realized as geodesic mapping of the equiaffine space $\tilde{A}_n$, which is projectively equivalent to the given space A_n .

2 Main properties of geodesic mappings

Let $A_n = (M, \nabla)$ be a manifold M with affine connection ∇ . We suppose that ∇ is torsion-free, i.e. $\nabla_X Y - \nabla_Y X = [X, Y]$, where $[,]$ is the Lie bracket: $[X, Y]f = X(Yf) - Y(Xf)$ for $f \in \mathcal{F}(M)$. Here and after $X, Y, \dots$ are tangent vectors.

The curvature R of a manifold A_n is a tensor field of type $(1, 3)$ defined by

$$R(X, Y)Z = \nabla_X(\nabla_Y Z) - \nabla_Y(\nabla_X Z) - \nabla_{[X, Y]}Z, \quad (1)$$

called sometimes also the Riemannian tensor of the connection.

We can introduce the Ricci tensor Ric of type $(0, 2)$ as a trace of a linear map, namely:

$$Ric(X, Y) = \text{trace}\{V \mapsto R(X, V)Y\}.$$

A manifold A_n with a torsion-free (symmetric) affine connection is called *equiaffine manifold*, if the Ricci tensor is symmetric, i.e. ([10, 11, 14])

$$Ric(X, Y) = Ric(Y, X).$$

It is known [10, 11, 14] that (pseudo-) Riemannian spaces are equiaffine manifolds.

A manifold $A_n = (M, \nabla)$ with affine connection admits a geodesic mapping onto $\bar{A}_n = (M, \bar{\nabla})$, if and only if the *Levi-Civita equation* holds [1, 3, 10, 14]:

$$\bar{\nabla}_X Y = \nabla_X Y + \psi(X)Y + \psi(Y)X, \quad (2)$$

where ∇ and $\bar{\nabla}$ are the affine connections of A_n and $\bar{A}_n$ respectively, and ψ is a linear form.

Geodesic mappings with $\psi \equiv 0$ are *trivial* or *affine*.

After applying formula (1) for the curvature tensor and expression (2) to a geodesic mapping $A_n \rightarrow \bar{A}_n$ we found a relationship between the curvature tensors R and $\bar{R}$ of A_n and $\bar{A}_n$:

$$\begin{aligned} \bar{R}(X, Y)Z &= R(X, Y)Z \\ &+ (\psi(X, Y) - \psi(Y, X))Z + \psi(X, Z)Y - \psi(Y, Z)X, \end{aligned} \quad (3)$$

where

$$\psi(X, Y) = \nabla_Y \psi(X) - \psi(X)\psi(Y). \quad (4)$$

By contraction of (3) we obtain the following relation for the Ricci tensors

$$\bar{Ric}(X, Y) = Ric(X, Y) + n\psi(X, Y) - \psi(Y, X). \quad (5)$$

3 Main results

Suppose $A_n \in C^1$, i.e. the components of the affine connection ∇ of A_n are functions of type C^1 on all charts of the manifold M .

In our paper [9] we proved that any manifold A_n with affine connection is locally projectively equiaffine.

The following theorem holds generally:

Theorem *All manifolds A_n with affine connection are projectively equiaffine.*

Remark. In other words, an arbitrary manifold $A_n \in C^1$ admits a global geodesic mapping onto an equiaffine manifold $\bar{A}_n$, and moreover $\bar{A}_n \in C^1$.

Proof. We prove the existence of a geodesic mapping of $A_n = (M, \nabla) \in C^1$ onto an equiaffine manifold $\bar{A}_n = (M, \bar{\nabla})$.

It is known that on the manifold M exists globally a metric, say $\tilde{g}$. Now we can construct the metric $\tilde{g}$ so that $\tilde{g} \in C^2$, i.e. the components $\tilde{g}_{ij}$ of $\tilde{g}$ in a coordinate domain of M are functions of type C^2 . We denote by $\tilde{\nabla}$ the Levi-Civita connection of $\tilde{g}$.

We construct the one-form ψ in the following way

$$\psi(X) = -\frac{1}{n+1} \text{trace}(Y \mapsto \nabla_X Y - \tilde{\nabla}_X Y). \quad (6)$$

With the help of formula (2) applied to ψ we construct globally the affine connection $\bar{\nabla}$ on M . It is evident that A_n is geodesically mapped onto $\bar{A}_n = (M, \bar{\nabla})$, and, evidently, $\bar{A}_n \in C^1$.

Now we prove that $\bar{A}_n$ is equiaffine. It is sufficient to calculate that

$$\bar{Ric}(X, Y) = \bar{Ric}(Y, X),$$

where $\bar{Ric}$ is the Ricci tensor of $\bar{A}_n$.

From (5) follows

$$Ric(X, Y) - Ric(Y, X) + (n + 1)(\psi(X, Y) - \psi(Y, X)) = 0,$$

where Ric is the Ricci tensor of A_n . Further we use relation (4):

$$Ric(X, Y) - Ric(Y, X) + (n + 1)(\nabla_Y \psi(X) - \nabla_X \psi(Y)) = 0.$$

After using (4) and (6) we can see that this equation holds identically. $\square$

Remark. The equiaffine connection $\bar{\nabla}$ constructed in this way is constructed explicitly from the original connection ∇ .

Similarly it could be shown that a manifold with projective connection admits a geodesic mapping onto a manifold with equiaffine connection.

References

- [1] M. Eastwood, V. Matveev, *Metric connections in projective differential geometry*, The IMA Volumes in Math. and its Appl. **144**, 339-350 (2008).
- [2] L.P. Eisenhart, *Riemannian Geometry*, Princeton Univ. Press, 1926.
- [3] L.P. Eisenhart, *Non-Riemannian geometry*, Amer. Math. Soc. Colloquium Publ. **8**, 2000.
- [4] S. Kobayashi and K. Nomizu, *Foundation of Differential Geometry* (Vol. 1, Interscience. Publ. New York–London, 1963; Vol. 2, Interscience. Publ., New York–London–Sydney, 1969).
- [5] T. Levi-Civita, *Sulle trasformazioni delle equazioni dinamiche*, Ann. Mat. Milano **24**, 2 (1886).
- [6] J. Mikeš, *Geodesic mappings of affine-connected and Riemannian spaces*, J. Math. Sci. **78**, 3 (1996).
- [7] J. Mikeš, V. Berezovski, *Geodesic mappings of affine-connected spaces onto Riemannian spaces*, Colloq. Math. Soc. J. Bolyai, **56**. Diff. Geom. Eger (Hungary), (1989).
- [8] J. Mikeš, I. Hinterleitner, V. Kiosak, *On the Theory of Geodesic Mappings of Einstein Spaces and their Generalizations*, AIP Conf. Proc. **861**, 428-435 (2006).

- [9] J. Mikeš, I. Hinterleitner, V. Kiosak, *On geodesic mappings of affine connection manifolds*, Acta Physica Debrecina **42**, 19-28 (2008).
- [10] J. Mikeš, V. Kiosak, A. Vanžurová, *Geodesic Mappings of Manifolds with Affine Connection*, Palacký University Press, 2008.
- [11] A.P. Norden, *Spaces of an Affine Connection*, Nauka, Moscow, 1976.
- [12] A.Z. Petrov, *New Methods in Theory of General Relativity*, Nauka, Moscow, 1966.
- [13] Ž. Radulović, J. Mikeš, M. Gavrilchenko, *Geodesic Mappings and Deformations of Riemannian Spaces*, Publisher, Podgorica, CID; Odessa, OGU, 1997.
- [14] N.S. Sinyukov *Geodesic Mappings of Riemannian Spaces*, Nauka, Moscow, 1979.
- [15] T.Y. Thomas, *On projective and equiprojective geometries of paths*, Proc. Nat. Acad. Sci. USA **11**, 198–203 (1925).